



acatalasemia

Acatalasemia is a condition characterized by very low levels of an enzyme called catalase. Many people with acatalasemia never have any health problems related to the condition and are diagnosed because they have affected family members.

Some of the first reported individuals with acatalasemia developed open sores (ulcers) inside the mouth that led to the death of soft tissue (gangrene). When mouth ulcers and gangrene occur with acatalasemia, the condition is known as Takahara disease. These complications are rarely seen in more recent cases of acatalasemia, probably because of improvements in oral hygiene.

Studies suggest that people with acatalasemia have an increased risk of developing type 2 diabetes mellitus, which is the most common form of diabetes. A higher percentage of people with acatalasemia have type 2 diabetes mellitus than in the general population, and the disease tends to develop at an earlier age (in a person's thirties or forties, on average). Researchers speculate that acatalasemia could also be a risk factor for other common, complex diseases; however, only a small number of cases have been studied.

Frequency

More than 100 cases of acatalasemia have been reported in the medical literature. Researchers estimate that the condition occurs in about 1 in 12,500 people in Japan, 1 in 20,000 people in Hungary, and 1 in 25,000 people in Switzerland. The prevalence of acatalasemia in other populations is unknown.

Genetic Changes

Mutations in the *CAT* gene can cause acatalasemia. This gene provides instructions for making the enzyme catalase, which breaks down hydrogen peroxide molecules into oxygen and water. Hydrogen peroxide is produced through chemical reactions within cells. At low levels, it is involved in several chemical signaling pathways, but at high levels it is toxic to cells. If hydrogen peroxide is not broken down by catalase, additional reactions convert it into compounds called reactive oxygen species that can damage DNA, proteins, and cell membranes.

Mutations in the *CAT* gene greatly reduce the activity of catalase. A shortage of this enzyme can allow hydrogen peroxide to build up to toxic levels in certain cells. For example, hydrogen peroxide produced by bacteria in the mouth may accumulate in and damage soft tissues, leading to mouth ulcers and gangrene. A buildup of hydrogen peroxide may also damage beta cells of the pancreas, which release a hormone called insulin that helps control blood sugar. Malfunctioning beta cells are thought to underlie

the increased risk of type 2 diabetes mellitus in people with acatalasemia. It is unclear why some people have no health problems associated with a loss of catalase activity.

Many people with reduced catalase activity do not have an identified mutation in the *CAT* gene; in these cases, the cause of the condition is unknown. Researchers believe that other genetic and environmental factors can also influence the activity of catalase.

Inheritance Pattern

Acatalasemia has an autosomal recessive pattern of inheritance, which means both copies of the *CAT* gene in each cell have mutations. When both copies of the gene are altered, the activity of catalase is reduced to less than 10 percent of normal.

When only one of the two copies of the *CAT* gene has a mutation, the activity of catalase is reduced by approximately half. This reduction in catalase activity is often called hypocatalasemia. Like acatalasemia, hypocatalasemia usually does not cause any health problems.

Other Names for This Condition

- acatalasia
- catalase deficiency

Diagnosis & Management

These resources address the diagnosis or management of acatalasemia:

- Genetic Testing Registry: Acatalasemia
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0268419/>
- Genetic Testing Registry: Acatalasemia, japanese type
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2936847/>

These resources from MedlinePlus offer information about the diagnosis and management of various health conditions:

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>

Additional Information & Resources

MedlinePlus

- Encyclopedia: Gangrene
<https://medlineplus.gov/ency/article/007218.htm>
- Encyclopedia: Mouth Ulcers
<https://medlineplus.gov/ency/article/001448.htm>
- Encyclopedia: Type 2 Diabetes
<https://medlineplus.gov/ency/article/000313.htm>
- Health Topic: Metabolic Disorders
<https://medlineplus.gov/metabolicdisorders.html>

Genetic and Rare Diseases Information Center

- Acatalasemia
<https://rarediseases.info.nih.gov/diseases/363/acatalasemia>

Additional NIH Resources

- National Institute of Diabetes and Digestive and Kidney Diseases: Your Guide to Diabetes: Type 1 and Type 2
<https://www.niddk.nih.gov/health-information/diabetes/overview/what-is-diabetes>

Educational Resources

- Children Living With Inherited Metabolic Diseases (CLIMB) (UK): Acatalasemia
<http://www.climb.org.uk/IMD/Alfa/Acatalasemia.pdf>
- Disease InfoSearch: Acatalasemia
<http://www.diseaseinfosearch.org/Acatalasemia/93>
- MalaCards: acatalasemia
<http://www.malacards.org/card/acatalasemia>
- Merck Manual Consumer Version: Diabetes Mellitus
<http://www.merckmanuals.com/home/hormonal-and-metabolic-disorders/diabetes-mellitus-dm-and-disorders-of-blood-sugar-metabolism/diabetes-mellitus-dm>
- Orphanet: Acatalasemia
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=926

Patient Support and Advocacy Resources

- American Diabetes Association
<http://www.diabetes.org/>
- Children Living With Inherited Metabolic Diseases (CLIMB) (UK)
<http://www.climb.org.uk/>

Genetic Testing Registry

- Acatalasemia
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0268419/>
- Acatalasemia, japanese type
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2936847/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28acatalasemia%5BMAJR%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D>

OMIM

- ACATALASEMIA
<http://omim.org/entry/614097>

Sources for This Summary

- Góth L, Eaton JW. Hereditary catalase deficiencies and increased risk of diabetes. *Lancet*. 2000 Nov 25;356(9244):1820-1.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11117918>
- Góth L, Nagy T. Acatalasemia and diabetes mellitus. *Arch Biochem Biophys*. 2012 Sep 15;525(2):195-200. doi: 10.1016/j.abb.2012.02.005. Epub 2012 Feb 16. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22365890>
- Góth L, Nagy T. Inherited catalase deficiency: is it benign or a factor in various age related disorders? *Mutat Res*. 2013 Oct-Dec;753(2):147-54. doi: 10.1016/j.mrrev.2013.08.002. Epub 2013 Sep 8. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24025477>
- Góth L, Rass P, Páy A. Catalase enzyme mutations and their association with diseases. *Mol Diagn*. 2004;8(3):141-9. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15771551>
- Góth L. A new type of inherited catalase deficiencies: its characterization and comparison to the Japanese and Swiss type of acatalasemia. *Blood Cells Mol Dis*. 2001 Mar-Apr;27(2):512-7.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11500062>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/acatalasemia>

Reviewed: September 2014

Published: February 7, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services